5

15

30

26

## Claims

- A transgenic non-human animal expressing at least one transgene comprising a DNA sequence encoding a heterologous Amyloid Precursor Protein (APP) comprising at least the Arctic mutation (E693G) and a further AD (Alzheimer's disease) pathogenic mutation or a further transgene affecting AD pathogenesis, which results in increased amounts of intracellular soluble Aβ aggregates, including Aβ peptides.
- 10 2. The transgenic animal according to claim 1, wherein the transgene/transgenes are integrated in the genomic DNA.
  - 3. The transgenic animal according to claim 1 or 2, wherein said transgene/transgenes are operably linked to a promoter effective for expression of said gene in the brain tissue of said animal.
    - 4. The transgenic animal according to any of claims 1-3 wherein the endogenous APP is expressive or non-expressive.
- 5. The transgenic animal according to any of claims 1-4, wherein said further transgene is a human presentiin-1 and/or presentiin-2 transgene harboring an AD pathogenic mutation.
- 6. The transgenic animal according to any of claims 1-4, wherein said further transgene comprises a DNA sequence encoding apolipoprotein E, apolipoprotein J (clusterin), a<sub>1</sub>-antichymotrypsin (ACT) or fragments thereof.
  - 7. The transgenic animal according to any of claims 1-4, wherein said further AD pathogenic mutation is one of the APP mutations KM670/671DF, KM670/671DY, KM670/671EF or KM670/671EY.
  - 8. The transgenic animal according to any of claims 1-4, wherein said further AD pathogenic mutation is one of the APP mutations KM670/671NL, KM670/671NY, KM670/671NF, KM670/671KL, KM670/671DL or KM670/671NDL and KM670/671NDL a
- 35 KM670/671EL, wherein KM670/671NL (the Swedish mutation) is preferred.

5

10

20

25

30

- 9. The transgenic animal according to any of claims 1-4, wherein the transgenic animal expresses only one transgene which comprises only the Arctic mutation (E693G) and the Swedish mutation (KM670/671NL).
- 10. The transgenic animal according to any of claims 1-9, additionally comprising a homologously integrated targeting construct for at least one of the neprilysin or insulin-degrading enzyme (IDE) genes, which disrupts these genes through gene ablation (knock-out) and enhances Aβ-40 and/or Aβ-42 Arctic peptide production.
- 11. The transgenic animal according to any of claims 1-10 wherein the transgenic animal is a rodent.
- 15 12. The transgenic animal according to any of claims 1-11 wherein the transgenic animal is a murine animal.
  - 13. The transgenic animal according to claim 1-12, wherein the transgenic animal is a mouse.
  - 14. A method of producing the transgenic animal according to any of claims 113, comprising integrating in the genomic DNA at least one transgene
    comprising a DNA sequence encoding a heterologous Amyloid Precursor Protein
    (APP) comprising at least the Arctic mutation (E693G) and a further AD
    (Alzheimer's disease) pathogenic mutation or a further transgene affecting AD
    pathogenesis.
  - 15. The method according to claim 14, wherein said transgene/transgenes are operably linked to a promoter effective for expression of said gene in the brain tissue of said animal.
  - 16. The method according to any of claims 14-15 wherein the endogenous APP is optionally made non-expressive.

20

25

- 17. The method according to any of claims 14-16, wherein said further transgene is a human presentiin-1 and/or presentiin-2 transgene harboring an AD pathogenic mutation.
- 5 18. The method according to any of claims 14-16, wherein said further transgene comprises a DNA sequence encoding apolipoprotein E, apolipoprotein J (clusterin), α<sub>1</sub>-antichymotrypsin (ACT) or fragments thereof.
- 19. The method according to any of claims 14, wherein said further AD pathogenic mutation is one of the APP mutations KM670/671DF, KM670/671DY, KM670/671EF or KM670/671EY.
  - 20. The method according to any of claims 14, wherein said further AD pathogenic mutation is one of the APP mutations KM670/671NL,
- 15 KM670/671NY, KM670/671NF, KM670/671KL, KM670/671DL or KM670/671EL, wherein KM670/671NL (the Swedish mutation) is preferred.
  - 21. The method according to any of claims 14-20, additionally comprising homologously integrating a targeting construct for at least one of the neprilysin or insulin-degrading enzyme (IDE) genes.
    - 22. A method of screening, wherein the transgenic animal according to any of claims 1-13 is used for screening for agents useful for treating, preventing or inhibiting Alzheimer's disease.
    - 23. A method of screening, wherein the transgenic animal according to any of claims 1-13 is used for screening for diagnostic agents for Alzheimer's disease.